

2017年度 協会けんぽ健診精密検査結果集計表（1月～12月）

| | 年 齢 | | —39 | | 40—44 | | 45—49 | | 50—54 | | 55—59 | | 60—64 | | 65—69 | | 70—74 | | 75—79 | | 80— | | |
|------------|-------|------|-------|-------|-------|------|-------|------|-------|-------|-------|------|-------|-------|-------|-------|-------|-------|-------|-----|-----|-----|-----|
| | 合計 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 |
| 胃X-P | | | | | | | | | | | | | | | | | | | | | | | |
| 受診者数 | 1,725 | 1045 | 680 | 187 | 103 | 208 | 151 | 183 | 137 | 141 | 115 | 126 | 85 | 102 | 63 | 78 | 22 | 18 | 4 | 2 | 0 | 0 | 0 |
| 要精検者数 | 86 | 63 | 23 | 5 | 1 | 8 | 4 | 8 | 2 | 5 | 4 | 14 | 7 | 10 | 4 | 11 | 1 | 2 | 0 | 0 | 0 | 0 | 0 |
| 要精検率 % | 5.0 | 6.0 | 3.4 | 2.7 | 1.0 | 3.8 | 2.6 | 4.4 | 1.5 | 3.5 | 3.5 | 11.1 | 8.2 | 9.8 | 6.3 | 14.1 | 4.5 | 11.1 | | | | | |
| 精検受診者数 | 45 | 29 | 16 | 2 | 0 | 3 | 4 | 3 | 1 | 4 | 1 | 6 | 6 | 3 | 3 | 6 | 1 | 2 | 0 | 0 | 0 | 0 | 0 |
| 精検受診率 % | 52.3 | 46.0 | 69.6 | 40.0 | 0.0 | 37.5 | 100.0 | 37.5 | 50.0 | 80.0 | 25.0 | 42.9 | 85.7 | 30.0 | 75.0 | 54.5 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 異常なし | 8 | 3 | 5 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 3 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 進行胃がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 早期胃がん | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 深達度不明胃がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 粘膜内がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃がん疑い | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃ポリープ | 5 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃潰瘍 | 4 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| 十二指腸潰瘍 | 6 | 5 | 1 | 1 | 0 | 0 | 1 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 共存潰瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 腺腫 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃粘膜下腫瘍 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 十二指腸ポリープ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 食道がん | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| 胃がん以外の悪性腫瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 不 明 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他 | 27 | 19 | 8 | 2 | 0 | 3 | 3 | 1 | 0 | 2 | 1 | 5 | 2 | 3 | 1 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃内視鏡検査 | | | | | | | | | | | | | | | | | | | | | | | |
| 受診者数 | 111 | 60 | 51 | 1 | 0 | 10 | 5 | 6 | 10 | 10 | 12 | 8 | 9 | 14 | 9 | 7 | 4 | 4 | 2 | 0 | 0 | 0 | 0 |
| 要精検者数 | 11 | 9 | 2 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 2 | 0 | 2 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| 要精検率 % | 9.9 | 15.0 | 3.9 | 100.0 | | 20.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.3 | 25.0 | 0.0 | 14.3 | 0.0 | 28.6 | 0.0 | 0.0 | 50.0 | | | | |
| 精検受診者数 | 10 | 8 | 2 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 2 | 0 | 2 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| 精検受診率 % | 90.9 | 88.9 | 100.0 | 100.0 | | 50.0 | | | | 100.0 | 100.0 | | | 100.0 | | 100.0 | | 100.0 | | | | | |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 異常なし | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 進行胃がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 早期胃がん | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 深達度不明胃がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 粘膜内がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃がん疑い | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃ポリープ | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃潰瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 十二指腸潰瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 共存潰瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 腺腫 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃粘膜下腫瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 十二指腸ポリープ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 食道がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃がん以外の悪性腫瘍 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 胃潰瘍癒痕 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 十二指腸潰瘍癒痕 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 不 明 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他 | 7 | 6 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |

| 胸部X-P | 年 齢 | | —39 | | 40—44 | | 45—49 | | 50—54 | | 55—59 | | 60—64 | | 65—69 | | 70—74 | | 75—79 | | 80— | | |
|------------|-------|-------|------|------|-------|-------|-------|------|-------|-------|-------|------|-------|-------|-------|------|-------|------|-------|-----|-----|-----|-----|
| | 合 計 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 |
| 受診者数 | 2,044 | 1,206 | 838 | 197 | 115 | 229 | 184 | 203 | 166 | 168 | 145 | 141 | 104 | 133 | 81 | 101 | 32 | 30 | 11 | 4 | 0 | 0 | 0 |
| 要精検者数 | 20 | 13 | 7 | 0 | 0 | 1 | 0 | 2 | 2 | 1 | 1 | 4 | 3 | 1 | 0 | 2 | 1 | 2 | 0 | 0 | 0 | 0 | 0 |
| 要精検率 % | 1.0 | 1.1 | 0.8 | 0.0 | 0.0 | 0.4 | 0.0 | 1.0 | 1.2 | 0.6 | 0.7 | 2.8 | 2.9 | 0.8 | 0.0 | 2.0 | 3.1 | 6.7 | 0.0 | 0.0 | | | |
| 精検受診者数 | 10 | 7 | 3 | 0 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 2 | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| 精検受診率 % | 50.0 | 53.8 | 42.9 | | | 100.0 | | 50.0 | 50.0 | 100.0 | 100.0 | 50.0 | 33.3 | 100.0 | | 0.0 | 0.0 | 50.0 | | | | | |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 肺がん疑い | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 肺結核 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 原発性肺がん0期 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 原発性肺がんⅠ期 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 原発性肺がんⅡ期 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 原発性肺がんⅢ期 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 原発性肺がんⅣ期 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他 | 3 | 2 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| 不 明 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 異常なし | 6 | 4 | 2 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他の肺悪性新生物 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他の肺新生物 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 便潜血 | | | | | | | | | | | | | | | | | | | | | | | |
| 受診者数 | 834 | 189 | 120 | 221 | 185 | 197 | 157 | 166 | 142 | 144 | 106 | 130 | 81 | 97 | 33 | 29 | 10 | 4 | 0 | 0 | 0 | 0 | 0 |
| 要精検者数 | 42 | 3 | 5 | 7 | 11 | 11 | 8 | 6 | 4 | 10 | 6 | 14 | 6 | 12 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 要精検率 % | 5.0 | 1.6 | 4.2 | 3.2 | 5.9 | 5.6 | 5.1 | 3.6 | 2.8 | 6.9 | 5.7 | 10.8 | 7.4 | 12.4 | 6.1 | 6.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 精検受診者数 | 17 | 3 | 0 | 3 | 6 | 5 | 3 | 2 | 2 | 7 | 2 | 3 | 2 | 3 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 精検受診率 % | 40.5 | 100.0 | 0.0 | 42.9 | 54.5 | 45.5 | 37.5 | 33.3 | 50.0 | 70.0 | 33.3 | 21.4 | 33.3 | 25.0 | 100.0 | 50.0 | | | | | | | |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 進行大腸がん | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 早期大腸がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 深達度不明大腸がん | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 大腸がんの疑い | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 大腸腺腫 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 2 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他大腸ポリープ | 2 | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 大腸憩室 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 潰瘍性大腸炎 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| クローン病 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| その他 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 不 明 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 異常なし | 11 | 3 | 0 | 2 | 5 | 3 | 2 | 1 | 2 | 2 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| P S A | 年 齢 | | —39 | | 40—44 | | 45—49 | | 50—54 | | 55—59 | | 60—64 | | 65—69 | | 70—74 | | 75—79 | | 80— | | |
|---------------|-------|------|-------|-----|-------|-----|-------|-------|-------|-------|-------|------|-------|------|-------|------|-------|-------|-------|-----|-----|---|---|
| | 合 計 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 |
| 受診者数 | 123 | 123 | | 5 | | 12 | | 13 | | 18 | | 23 | | 17 | | 24 | | 10 | | 1 | | 0 | |
| 要精検者数 | 11 | 11 | | 0 | | 1 | | 1 | | 1 | | 3 | | 2 | | 2 | | 1 | | 0 | | 0 | |
| 要精検率 % | 8.9 | 8.9 | | 0.0 | | 8.3 | | 7.7 | | 5.6 | | 13.0 | | 11.8 | | 8.3 | | 10.0 | | 0.0 | | | |
| 精検受診者数 | 6 | 6 | | 0 | | 0 | | 1 | | 1 | | 1 | | 1 | | 1 | | 1 | | 0 | | 0 | |
| 精検受診率 % | 54.5 | 54.5 | | | | 0.0 | | 100.0 | | 100.0 | | 33.3 | | 50.0 | | 50.0 | | 100.0 | | | | | |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 進行がん | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 早期がんB 0 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 早期がんB 1 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 早期がんB 2 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 局所進展がん | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| がんの疑い | 3 | 3 | | 0 | | 0 | | 0 | | 0 | | 1 | | 0 | | 1 | | 1 | | 0 | | 0 | |
| 病気不明がん | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| その他 | 1 | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 | |
| 不 明 | 0 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 異常なし | 2 | 2 | | 0 | | 0 | | 1 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | |
| 乳 | | | | | | | | | | | | | | | | | | | | | | | |
| 受診者数 | 362 | | 362 | | 24 | | 101 | | 57 | | 77 | | 46 | | 38 | | 13 | | 6 | | 0 | | 0 |
| 要精検者数 | 27 | | 27 | | 1 | | 9 | | 5 | | 7 | | 1 | | 2 | | 1 | | 1 | | 0 | | 0 |
| 要精検率 % | 7.5 | | 7.5 | | 4.2 | | 8.9 | | 8.8 | | 9.1 | | 2.2 | | 5.3 | | 7.7 | | 16.7 | | | | |
| 精検受診者数 | 14 | | 14 | | 1 | | 7 | | 4 | | 1 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 |
| 精検受診率 % | 51.9 | | 51.9 | | 100.0 | | 77.8 | | 80.0 | | 14.3 | | 0.0 | | 50.0 | | 0.0 | | 0.0 | | | | |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 乳がん | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| がんの疑い | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 線維腺腫 | 3 | | 3 | | 0 | | 0 | | 2 | | 0 | | 0 | | 0 | | 1 | | 0 | | 0 | | 0 |
| 乳腺症 | 4 | | 4 | | 0 | | 1 | | 1 | | 2 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| のう胞 | 6 | | 6 | | 0 | | 3 | | 2 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 良性石灰化像 | 3 | | 3 | | 0 | | 1 | | 1 | | 0 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 |
| その他 | 4 | | 4 | | 1 | | 1 | | 1 | | 0 | | 0 | | 0 | | 0 | | 1 | | 0 | | 0 |
| 不 明 | 1 | | 1 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 異常なし | 6 | | 6 | | 0 | | 1 | | 0 | | 4 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 |
| ※要精検数はMMG結果のみ | | | | | | | | | | | | | | | | | | | | | | | |
| 子宮 | | | | | | | | | | | | | | | | | | | | | | | |
| 受診者数 | 361 | | 361 | | 50 | | 90 | | 49 | | 79 | | 41 | | 36 | | 11 | | 5 | | 0 | | 0 |
| 要精検者数 | 15 | | 15 | | 5 | | 6 | | 2 | | 2 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 要精検率 % | 4.2 | | 4.2 | | 10.0 | | 6.7 | | 4.1 | | 2.5 | | 0.0 | | 0.0 | | 0.0 | | 0.0 | | | | |
| 精検受診者数 | 15 | | 15 | | 5 | | 6 | | 2 | | 2 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 精検受診率 % | 100.0 | | 100.0 | | 100.0 | | 100.0 | | 100.0 | | 100.0 | | | | | | | | | | | | |
| 結 果 | | | | | | | | | | | | | | | | | | | | | | | |
| 浸潤がん(子宮頸) | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 上皮内がん(子宮頸) | 3 | | 3 | | 1 | | 1 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 子宮体がん | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 高度異形成 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 中等度異形成 | 1 | | 1 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 軽度異形成 | 2 | | 2 | | 1 | | 0 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 内膜増殖症 | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| その他 | 1 | | 1 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 不 明 | 1 | | 1 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| 異常なし | 3 | | 3 | | 1 | | 0 | | 1 | | 1 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 |
| ※要精検数は細胞診結果のみ | | | | | | | | | | | | | | | | | | | | | | | |